

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number  
**WO 01/35961 A1**

(51) International Patent Classification?: **A61K 31/535**,  
31/50, 31/495, 31/415, 31/355, 31/34, 31/19, 31/135,  
A01N 45/00

**WORCEL, Manuel** [FR/US]; 20 Gloucester Street, No.  
4, Boston. MA 02115 (US).

(21) International Application Number: **PCT/US00/29528**

(22) International Filing Date: 27 October 2000 (27.10.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/162,230 29 October 1999 (29.10.1999) US  
60/179,020 31 January 2000 (31.01.2000) US

(74) Agent: **GRIEFF, Edward, D.**; Hale and Dorr LLP, 1455  
Pennsylvania Avenue, N.W., Washington, D C20004 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants (*for all designated States except US*): **NI-  
TROMED, INC.** [US/US]; 12 Oak Park Drive, Bedford,  
MA 01730 (US). **TRUSTEES OF BOSTON UNIVER-  
SITY** [US/US]; 147 Bay State Road, Boston, MA 02215  
(US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **LOSCALZO**,  
**Joseph** [US/US]; 16 LedgeWood Drive, Dover, MA 02030  
(US). **VITA, Joseph, A.** [US/US]; 12 Fulling Mill Lane,  
Hingham, MA 02118 (US). **LOBERG, Michael, D.**  
[US/US]; 82 Beacon Street, Boston, MA 02116 (US).

**Published:**

- With international search report.
- Before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments.

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: METHODS OF TREATING VASCULAR DISEASES CHARACTERIZED BY NITRIC OXIDE INSUFFICIENCY

(57) Abstract: The present invention provides methods of treating and/or preventing vascular diseases where nitric oxide insufficiency is a contributing factor by administering a therapeutically effective amount of at least one antioxidant, or a pharmaceutically acceptable salt thereof, and at least one of isosorbide dinitrate and isosorbide mononitrate, and, optionally, at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, and/or at least one compound used to treat cardiovascular diseases. The antioxidant is preferably a hydralazine compound or a pharmaceutically acceptable salt thereof. The present invention also provides methods of treating and/or preventing vascular diseases where nitric oxide insufficiency is a contributing factor by administering a therapeutically effective amount of at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated angiotensin II receptor antagonist, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, and/or at least one compound used to treat cardiovascular diseases. The present invention also provides methods of treating and/or preventing Raynaud's syndrome by administering a therapeutically effective amount of at least one antioxidant and at least one of isosorbide dinitrate and isosorbide mononitrate and/or at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor. The present invention also provides novel transdermal patches comprising at least one antioxidant and at least one of isosorbide dinitrate and isosorbide mononitrate and/or at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor.

WO 01/35961 A1

**METHODS OF TREATING VASCULAR DISEASES CHARACTERIZED BY  
NITRIC OXIDE INSUFFICIENCY  
RELATED APPLICATIONS**

This application claims priority to U. S. Provisional Application No.  
5 60/162,230 filed October 29, 1999 and U. S. Provisional Application No. 60/179,020  
filed January 31, 2000.

**FIELD OF THE INVENTION**

The present invention provides methods of treating and/or preventing  
vascular diseases characterized by nitric oxide insufficiency by administering a  
therapeutically effective amount of at least one antioxidant or a pharmaceutically  
10 acceptable salt thereof, and at least one of isosorbide dinitrate and isosorbide  
mononitrate, and, optionally, at least one nitrosated angiotensin-converting enzyme  
inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker,  
nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist,  
15 nitrosated renin inhibitor, and/or at least one compound used to treat  
cardiovascular diseases. The antioxidant is preferably a hydralazine compound or a  
pharmaceutically acceptable salt thereof. The present invention also provides  
methods of treating and/or preventing vascular diseases characterized by nitric  
oxide insufficiency by administering a therapeutically effective amount of at least  
20 one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic  
blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist,  
nitrosated angiotensin II receptor antagonist and/or nitrosated renin inhibitor, and,  
optionally, at least one antioxidant and/or at least one compound used to treat  
cardiovascular diseases. The present invention also provides methods of treating  
25 and/or preventing Raynaud's syndrome by administering a therapeutically effective  
amount of at least one antioxidant or a pharmaceutically acceptable salt thereof, and  
at least one of isosorbide dinitrate and isosorbide mononitrate, and/or at least one  
nitrosated angiotensin-converting enzyme inhibitor, nitrosated calcium channel  
blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor  
30 antagonist and/or nitrosated renin inhibitor. The present invention also provides  
novel transdermal patches comprising at least one antioxidant or a pharmaceutically  
acceptable salt thereof, and at least one of isosorbide dinitrate and isosorbide  
mononitrate, and/or at least one nitrosated angiotensin-converting enzyme

inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist and/or nitrosated renin inhibitor.

### BACKGROUND OF THE INVENTION

5       The decline in cardiovascular morbidity and mortality in the United States over the past three decades has been the result of significant advances in research on cardiovascular disease mechanisms and therapeutic strategies. The incidence and prevalence of myocardial infarction and death from myocardial infarction, as well as  
10       largely owing to advances in prevention, early diagnosis, and treatment of these very common diseases.

      Analysis of outcomes by race, however, paints quite a different picture: life expectancy and cardiovascular morbidity rates have improved far less for blacks than whites. Available data show that the likelihood of dying from cardiovascular  
15       disease is far greater among black Americans than among white Americans. In this decade, the death rate from cardiovascular disease for black males was 353 per 100,000 population, while that for white males was 244 per 100,000; the rate for black females was 226 per 100,000; while that for white females was 135 per 100,000. Consonant with this important demographic parameter is the observation that there  
20       is a higher prevalence of several of the important risk factors for cardiovascular disease, e.g., hypertension, smoking, diabetes mellitus, obesity, and left ventricular hypertrophy, among blacks compared with whites. In addition, outcomes of cardiovascular events are worse for blacks than whites. Following myocardial infarction, blacks have a 50% higher annual mortality rate than whites, and their five  
25       year survival is only 70%. Thus, the many advances in cardiovascular medicine that account for the overall improvement in cardiovascular health in the general population have failed to translate into comparable racial benefits.

      There is a need in the art for new and more effective compositions and methods for treating vascular diseases. The present invention is directed to these, as  
30       well as other, important ends.

### SUMMARY OF THE INVENTION

      The present invention provides methods for treating and/or preventing vascular diseases characterized by nitric oxide insufficiency by administering to a

patient a therapeutically effective amount of at least one antioxidant or a pharmaceutically acceptable salt thereof; and at least one of isosorbide dinitrate and isosorbide mononitrate, and, optionally, at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, and/or at least one compound used to treat cardiovascular diseases. The antioxidant is preferably a hydralazine compound or a pharmaceutically acceptable salt thereof. The vascular diseases characterized by nitric oxide insufficiency include, for example, hypertension (e.g., low-renin hypertension; salt-sensitive hypertension; low-renin, salt-sensitive hypertension; primary pulmonary hypertension; thromboembolic pulmonary hypertension; pregnancy-induced hypertension; renovascular hypertension), heart failure (e.g., microvascular cardiac ischemia), and left ventricular hypertrophy with disproportionate microvascularization, (i.e., inadequate vascularity) or diastolic dysfunction. The antioxidant and the isosorbide dinitrate or isosorbide mononitrate and optional nitrosated compound and/or compound used to treat cardiovascular diseases can be administered separately or as components of the same composition.

Another aspect of the present invention provides methods for treating and/or preventing vascular diseases characterized by nitric oxide insufficiency by administering to a patient a therapeutically effective amount of at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist and/or nitrosated renin inhibitor, and, optionally, at least one antioxidant and/or at least one compound used to treat cardiovascular diseases. The vascular diseases characterized by nitric oxide insufficiency include, for example, hypertension (e.g., low-renin hypertension; salt-sensitive hypertension; low-renin, salt-sensitive hypertension; primary pulmonary hypertension; thromboembolic pulmonary hypertension; pregnancy-induced hypertension; renovascular hypertension), heart failure (e.g., microvascular cardiac ischemia), and left ventricular hypertrophy with disproportionate microvascularization, (i.e., inadequate vascularity) or diastolic dysfunction. The nitrosated compound and optional antioxidant and/or compound used to treat cardiovascular diseases can be administered separately or as components of the

same composition.

In another aspect, the present invention provides methods for treating and/or preventing Raynaud's syndrome by administering to a patient a therapeutically effective amount of at least one antioxidant or a pharmaceutically acceptable salt thereof, and at least one of isosorbide dinitrate and isosorbide mononitrate and/or at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist and/or nitrosated renin inhibitor. The antioxidant, isosorbide dinitrate or isosorbide mononitrate, and nitrosated compound can be administered separately or as components of the same composition.

In yet another aspect, the present invention provides novel transdermal patches comprising a therapeutically effective amount of at least one antioxidant at least one of isosorbide dinitrate and isosorbide mononitrate and/or at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist and/or nitrosated renin inhibitor, and mixtures thereof.

These and other aspects of the present invention are described in more detail herein.

#### BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 shows that nitric oxide (NO) insufficiency is associated with increased salt and water retention and a low-renin state. Increased intracellular sodium and calcium in conjunction with reduced NO leads to enhanced sensitivity of vascular smooth muscle cells and cardiomyocytes to the tonic and growth-stimulating properties of catecholamines and angiotensin II. Increased vascular tone, left ventricular hypertrophy with inadequate capillary angiogenesis, and increased matrix production with myocardial fibrosis result. These intermediate phenotypes lead to the clinical disorders of low-renin, salt-sensitive hypertension; disproportionate left ventricular hypertrophy and diastolic dysfunction; and microvascular myocardial ischemia.

Fig. 2 shows forearm blood flow responses to intra-arterial methacholine that were assessed using venous occlusion plethysmography in 20 white and 16 black patients with a clinical history of hypertension (BP>140/90). Fig. 2 shows that

endothelium-derived NO action is impaired in the forearm microvessels of the black patients compared to the white patients.

5 Figs. 3A and 3B show forearm blood flow responses to nitroprusside (Fig. 3A) and methacholine (Fig. 3B) that were assessed by venous occlusion plethysmography in 25 white and 21 black patients without hypertension. The dilator response to sodium nitroprusside (Fig. 3A) was significantly lower in black patients, while there was no racial difference in response to methacholine (Fig. 3B).

10 Figs. 4A and 4B show the effect of salt-sensitivity on forearm microvascular function. By repeated measures ANOVA, there were trends for impaired responses to methacholine (Fig. 4A) and sodium nitroprusside (Fig. 4B) in salt-sensitive black patients.

### DETAILED DESCRIPTION OF THE INVENTION

As used throughout the disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings.

15 "Patient" refers to animals, preferably mammals, most preferably humans, and includes males and females.

"Therapeutically effective amount" refers to the amount of the compound and/or composition that is effective to achieve its intended purpose.


20 "Transdermal" refers to the delivery of a compound by passage through the skin and into the blood stream.

"Transmucosal" refers to delivery of a compound by passage of the compound through the mucosal tissue and into the blood stream.

25 "Penetration enhancement" or "permeation enhancement" refers to an increase in the permeability of the skin or mucosal tissue to a selected pharmacologically active compound such that the rate at which the compound permeates through the skin or mucosal tissue is increased.

30 "Carriers" or "vehicles" refers to carrier materials suitable for compound administration and include any such material known in the art such as, for example, any liquid, gel, solvent, liquid diluent, solubilizer, or the like, which is non-toxic and which does not interact with any components of the composition in a deleterious manner.

"Hydralazine compound" refers to a compound having the formula:



wherein a, b and c are independently a single or double bond; R<sub>1</sub> and R<sub>2</sub> are each independently a hydrogen, an alkyl, an ester or a heterocyclic ring; R<sub>3</sub> and R<sub>4</sub> are each independently a lone pair of electrons or a hydrogen, with the proviso that at least one of R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> is not a hydrogen. Exemplary hydralazine compounds include budralazine, cadralazine, dihydralazine, endralazine, hydralazine, pildralazine, todralazine, and the like.

"Antioxidant" refers to a compound that can react and quench a free radical.

"Alkyl" refers to a lower alkyl group, a haloalkyl group, a hydroxyalkyl group, an alkenyl group, an alkynyl group, a bridged cycloalkyl group, a cycloalkyl group or a heterocyclic ring, as defined herein.

"Lower alkyl" refers to branched or straight chain acyclic alkyl group comprising one to about ten carbon atoms (preferably one to about eight carbon atoms, more preferably one to about six carbon atoms). Exemplary lower alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, pentyl, neopentyl, iso-amyl, hexyl, octyl, and the like.

"Haloalkyl" refers to a lower alkyl group, an alkenyl group, an alkynyl group, a bridged cycloalkyl group, a cycloalkyl group or a heterocyclic ring, as defined herein, to which is appended one or more halogens, as defined herein. Exemplary haloalkyl groups include trifluoromethyl, chloromethyl, 2-bromobutyl, 1-bromo-2-chloro-pentyl, and the like.

"Alkenyl" refers to a branched or straight chain C<sub>2</sub>-C<sub>10</sub> hydrocarbon (preferably a C<sub>2</sub>-C<sub>8</sub> hydrocarbon, more preferably a C<sub>2</sub>-C<sub>6</sub> hydrocarbon) which can comprise one or more carbon-carbon double bonds. Exemplary alkenyl groups include propylenyl, buten-1-yl, isobutenyl, penten-1-yl, 2,2-methylbuten-1-yl, 3-methylbuten-1-yl, hexan-1-yl, hepten-1-yl, octen-1-yl, and the like.

"Alkynyl" refers to an unsaturated acyclic C<sub>2</sub>-C<sub>10</sub> hydrocarbon (preferably a C<sub>2</sub>-C<sub>8</sub> hydrocarbon, more preferably a C<sub>2</sub>-C<sub>6</sub> hydrocarbon) which can comprise one or more carbon-carbon triple bonds. Exemplary alkynyl groups include ethynyl, propynyl, butyn-1-yl, butyn-2-yl, pentyl-1-yl, pentyl-2-yl, 3-methylbutyn-1-yl, hexyl-

1-yl, hexyl-2-yl, hexyl-3-yl, 3,3-dimethyl-butyn-1-yl, and the like.

"Bridged cycloalkyl" refers to two or more cycloalkyl groups, heterocyclic groups, or a combination thereof fused via adjacent or non-adjacent atoms. Bridged cycloalkyl groups can be unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, amino, alkylamino, dialkylamino, hydroxy, halo, carboxyl, alkylcarboxylic acid, aryl, amidyl, ester, alkylcarboxylic ester, carboxamido, alkylcarboxamido, oxo and nitro. Exemplary bridged cycloalkyl groups include adamantyl, decahydronaphthyl, quinuclidyl, 2,6-dioxabicyclo[3.3.0]octane, 7-oxabicyclo[2.2.1]heptyl, 8-azabicyclo[3.2.1]oct-2-enyl, and the like.

"Cycloalkyl" refers to a saturated or unsaturated cyclic hydrocarbon comprising from about 3 to about 8 carbon atoms. Cycloalkyl groups can be unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, amino, alkylamino, dialkylamino, arylamino, diarylamino, alkylaryl amino, aryl, amidyl, ester, hydroxy, halo, carboxyl, alkylcarboxylic acid, alkylcarboxylic ester, carboxamido, alkylcarboxamido, oxo and nitro. Exemplary cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, cyclohepta,1,3-dienyl, and the like.

"Heterocyclic ring or group" refers to a saturated or unsaturated cyclic hydrocarbon group having about 2 to about 10 carbon atoms (preferably about 4 to about 6 carbon atoms) where 1 to about 4 carbon atoms are replaced by one or more nitrogen, oxygen and/or sulfur atoms. Sulfur maybe in the thio, sulfinyl or sulfonyl oxidation state. The heterocyclic ring or group can be fused to an aromatic hydrocarbon group. Heterocyclic groups can be unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, amino, alkylamino, dialkylamino, arylamino, diarylamino, alkylaryl amino, hydroxy, oxo, thial, halo, carboxyl, carboxylic ester, alkylcarboxylic acid, alkylcarboxylic ester, aryl, arylcarboxylic acid, arylcarboxylic ester, alkylcarbonyl, arylcarbonyl, amidyl, ester, carboxamido, alkylcarboxamido, arylcarboxamido, hydrazino, sulfonic acid, sulfonic ester, sulfonamido and nitro. Exemplary heterocyclic groups include pyrrolyl, 3-pyrrolinyl, 4,5,6-trihydro-2H-pyranyl, pyridinyl, 1,4-dihydropyridinyl, pyrazolyl, triazolyl, pyrimidinyl, pyridazinyl, oxazolyl, thiazolyl, imidazolyl, indolyl, thiophenyl, furanyl, tetrahydrofuranyl, tetrazolyl, 2-pyrrolinyl, 3-pyrrolinyl,



pyrrolindinyl, oxazolindinyl, 1,3-dioxolanyl, 2-imidazonlinyl, imidazolindinyl, 2-pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, 1,2,3-oxadiazolyl, 1,2,3-triazolyl, 1,3,4-thiadiazolyl, 2H-pyranyl, 4H-pyranyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl, 1,3,5-trithianyl, benzo(b)thiophenyl, benzimidazolyl, quinolinyl, and the like.

"Aryl" refers to a monocyclic, bicyclic, carbocyclic or heterocyclic ring system comprising one or two aromatic rings. Exemplary aryl groups include phenyl, pyridyl, naphthyl, quinoyl, tetrahydronaphthyl, furanyl, indanyl, indenyl, indoyl, and the like. Aryl groups (including bicyclic aryl groups) can be unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, amino, alkylamino, dialkylamino, arylamino, diarylamino, alkylaryl amino, hydroxy, carboxyl, carboxylic ester, alkylcarboxylic acid, alkylcarboxylic ester, aryl, arylcarboxylic acid, arylcarboxylic ester, alkylcarbonyl, arylcarbonyl, amidyl, ester, carboxamido, alkylcarboxamido, carbonyl, sulfonic acid, sulfonic ester, sulfonamido and nitro. Exemplary substituted aryl groups include tetrafluorophenyl, pentafluorophenyl, sulfonamide, alkylsulfonyl, arylsulfonyl, and the like.

"Hydroxy" refers to -OH.

"Hydroxyalkyl" refers to a hydroxy group, as defined herein, appended to an alkyl group, as defined herein.

"Alkylcarbonyl" refers to  $R_{52}-C(O)-$ , wherein  $R_{52}$  is an alkyl group, as defined herein.

"Arylcarbonyl" refers to  $R_{55}-C(O)-$ , wherein  $R_{55}$  is an aryl group, as defined herein.

"Ester" refers to  $R_{51}C(O)O-$  wherein  $R_{51}$  is a hydrogen atom, an alkyl group, an aryl group, an alkylaryl group, or an arylheterocyclic ring, as defined herein.

"Alkylaryl" refers to an alkyl group, as defined herein, to which is appended an aryl group, as defined herein. Exemplary alkylaryl groups include benzyl, phenylethyl, hydroxybenzyl, fluorobenzyl, fluorophenylethyl, and the like.

"Arylheterocyclic ring" refers to a bi- or tricyclic ring comprised of an aryl ring, as defined herein, appended via two adjacent carbon atoms of the aryl ring to a heterocyclic ring, as defined herein. Exemplary arylheterocyclic rings include dihydroindole, 1,2,3,4-tetra-hydroquinoline, and the like.

"Hydrazino" refers to  $H_2N-N(H)-$ .

“Compound used to treat cardiovascular diseases” refers to any therapeutic compound, or a pharmaceutically acceptable salt thereof, used to treat any cardiovascular disease. Suitable compounds include, but are not limited to, angiotensin-converting enzyme (ACE) inhibitors (such as, for example, alacepril, benazepril, captopril, ceranapril, cilazapril, delapril, duinapril, enalapril, enalaprilat, fosi-  
5 nopril, imidapril, lisinopril, moveltipril, pentopril, perindopril, quinapril, ramipril, rentipril, spirapril, temocapril,trandolapril, zofenopril, and the like); beta-adrenergic blockers (such as, for example, amosulalol, atenolol, betaxolol, bethanidine, bevantolol, bisoprolol, bopindolol, bufuralol, bunitrolol, bupranolol, buta-  
10 filolol, carazolol, carteolol, carvedilol, celiprolol, cetamolol, dilevalol, epanolol, indenolol, mepindolol, metipranolol, metoprolol, moprolol, nadolol, nebivolol, nipradilol, penbutolol, pindolol, pronethalol, propranolol, sotalol, sulfinalol, talinolol, tertatolol, tilisolol, timolol, toliprolol, and the like); cholesterol reducers (such as, for example, HMG-CoA reductase inhibitors, including, but not limited to, lovastatin (MEVACOR®), simvastatin (ZOCOR®), pravastatin (PRAVACHOL®),  
15 fluvastatin, cerivastatin (BAYCOL®), atorvastatin (LIPITOR®), and the like; sequestrants, including, but not limited to, cholestyramine, colestipol, sialkylaminoalkyl derivatives of cross-linked dextran, and the like; inhibitors of cholesterol absorption, including, but not limited to, beta-sitosterol, acyl CoA-  
20 cholersterol acyltransferase inhibitors, melinamide, and the like); calcium channel blockers (such as, for example, amlodipine, aranidipine, barnidipine, benidipine, cilnidipine, clentiazem, diltiazem, efonidipine, fantofarone, felodipine, isradipine, lacidipine, lercanidipine, manidipine, mibefradil, nicardipine, nifedipine, nilvadipine, nisoldipine, nitrendipine, semotiadil, veraparmil, and the like);  
25 angiotensin II receptor antagonists (such as, for example, ciclosidomine, eprosartan, furosemide, irbesartan, losartan, saralasin, valsartan, and the like); endothelin antagonists (such as, for example, bosentan, sulfonamide endothelin antagonists, BQ-123, SQ 28608, and the like); renin inhibitors (such as, for example, enalkrein, RO 42-5892, A 65317, CP 80794, ES 1005, ES 8891, SQ 34017, and the like); and mixtures  
30 thereof.

“Cardiovascular diseases” refers to any cardiovascular disease, including but not limited to, congestive heart failure, hypertension, pulmonary hypertension, myocardial and cerebral infarctions, atherosclerosis, atherogenesis, thrombosis,

ischemic heart disease, post-angioplasty restenosis, coronary artery diseases, renal failure, stable, unstable and variant (Prinzmetal) angina, cardiac edema, renal insufficiency, nephrotic edema, hepatic edema, stroke, and the like.

Two broad classes of cardiovascular disorders are more prevalent among blacks than whites and serve as areas in need of investigative efforts. Hypertension and left ventricular hypertrophy, two related yet independent risk factors for coronary heart disease, are significantly more prevalent among blacks than whites. Blacks also have higher rates of angiographically normal coronary arteries despite a higher prevalence of risk factors for coronary atherosclerosis, and greater morbidity and mortality from coronary heart disease than whites. These paradoxical observations have led some investigators to postulate that blacks harbor a diathesis of the microvasculature that limits perfusion and serves as a stimulus for vascular smooth muscle cell and cardiomyocyte hypertrophy, which, in turn, leads to hypertension and left ventricular hypertrophy, respectively. The underlying basis for this vascular diathesis may involve the endothelium, which has a limited capacity to generate vasodilator and antiproliferative factors or an increased capacity to produce vasoconstrictor and proliferative factors; the vascular smooth muscle cell, which manifests increased sensitivity to vasoconstrictor and proliferative factors; or both, in these individuals.

The present inventors have discovered that a major product of the normal blood vessel that may play a role in the vascular diathesis of blacks is endothelium-derived nitric oxide (NO). Nitric oxide produced by the endothelial cells induces vascular smooth muscle cell relaxation, contributing importantly to resting vascular tone. In addition, NO inhibits vascular smooth muscle cell proliferation and induces apoptosis in smooth muscle cells, which leads to the release of basic fibroblast growth factor and vascular endothelial cell growth factor, in turn supporting endothelial cell proliferation. This sequence of cellular responses is believed to sustain angiogenesis under hypoxic or ischemic conditions.

The role of NO in the vascular diathesis of blacks is illustrated by the consequences of NO insufficiency in the normal responses of the vasculature to NO. Nitric oxide insufficiency suppresses renin release from the juxtaglomerular cells, and induces a sodium chloride/volume sensitive increase in blood pressure. Furthermore, NO insufficiency leads to an increased sensitivity of vascular smooth

muscle cells to vasoconstrictors, such as angiotensin II and catecholamines, which amplify the increase in vascular resistance.

Nitric oxide insufficiency promotes vascular smooth muscle cell proliferation following vascular injury, and sustains smooth muscle cell and cardiomyocyte hypertrophy in response to catecholamines and angiotensin II. Furthermore, inadequate NO leads to increased production of extracellular matrix with consequent myocardial fibrosis.

These many cardiovascular responses that result from inadequate NO in the vasculature have clear clinical correlates in the black population. The clinical vascular phenotype of blacks that distinguishes them from whites with similar cardiovascular disorders is one of salt-sensitive, low-renin hypertension; left ventricular hypertrophy disproportionate to afterload and with an inadequate angiogenic response; and microvascular ischemia in the absence of significant epicardial coronary artery disease. The net pathophysiological consequences of these effects are increased peripheral vascular resistance with accompanying arterial hypertension; and an inadequately vascularized, fibrotic increase in left ventricular mass with accompanying diastolic dysfunction and microvascular ischemia.

Given these clinical observations and the role that NO plays in preventing their development, the present inventors have unexpectedly discovered that the principal cardiovascular disorders common among blacks (such as hypertension, left ventricular hypertrophy, and heart failure) result from a specific vascular diathesis that is a direct consequence of nitric oxide insufficiency. An outline of the pathogenic consequences of NO insufficiency that serve as the basis for these cardiovascular disorders is shown in Fig. 1.

NO insufficiency states can be a consequence of reduced synthesis of NO, enhanced inactivation of NO, or both. Possible candidate mechanisms include alterations in the genes that code for endothelial NO synthase or the inducible microvascular and cardiomyocyte NO synthase leading to reduced expression of a normal gene product or appropriate expression of a less active gene product; reduction in the enzymatic activity of NO synthase owing to inadequate cofactor concentrations; or enhanced inactivation of NO by oxidant stress.

Data obtained by the inventors in cultured cells, animal models, and human patients suggest that increased oxidant stress is central to the vascular diathesis of

and consequent cardiovascular disorders common among African Americans. Possible candidate mechanisms for the oxidant stress include enhanced production of reactive oxygen species (ROS), decreased antioxidant defenses, or both. The inventors make no *a priori* assumptions about the temporal or causative relationship between oxidant stress and the vascular phenotype of blacks: oxidant stress may both precede the development of the vascular diathesis and promote its progression once established. Recent data suggest that enhanced ROS production accompanies essential hypertension, atherosclerosis, thrombosis, and diabetes mellitus, and appears in each case, at the very least, to be important in the progression of established disease, if not in its actual genesis.

Endothelium-derived relaxing factor (EDRF), first described by Furchgott et al, *Nature*, 299:373-376 (1980), is an important mediator of vascular function. This endothelial product activates guanylyl cyclase in vascular smooth muscle cells and platelets, leading to vasorelaxation and platelet inhibition, respectively (Loscalzo et al, *Prog Cardiovasc Dis*, 38:87-104 (1995)). The chemical nature of EDRF has been studied using a variety of pharmacological and analytical techniques, and is NO (Ignarro et al, *Circ Res*, 61:866-879 (1987); Palmer et al, *Nature*, 327:524-526 (1987)).

Nitric oxide is synthesized by one of several isoforms of the NO synthase (NOS) family of enzymes, two of which are found in the vasculature, endothelial NOS (eNOS) and inducible NOS (iNOS). eNOS is synthesized by endothelial cells, while iNOS is synthesized by a variety of cell types, including vascular smooth muscle cells, fibroblasts, and (principally microvascular) endothelial cells (Balligand et al, *Am J Physiol*, 268:H1293-1303 (1995)). These enzymes produce NO as a result of the five-electron oxidation of L-arginine to L-citrulline; requisite cofactors include calcium-calmodulin, O<sub>2</sub>, FAD, FMN, tetrahydrobiopterin thiols, heme, and NADPH. (Moncada et al, *N Engl J Med*, 329:2002-2012 (1993)).

The role of NO in the cardiovascular system has become increasingly apparent over the past fifteen years (Loscalzo et al, *Prog Cardiovasc Dis*, 38:87-104 (1995)). Nitric oxide contributes importantly to resting tone in conductance as well as resistance arteries (Ouyyumi et al, *J Clin Invest*, 95:1747-1755 (1995)), and plays a critical role in the maintenance of peripheral vascular resistance and arterial pressure responses. Inhibition of NOS activity is associated with enhanced vascular sensitivity to vasoconstrictors, such as norepinephrine and angiotensin II (Conrad et

al, *Am J Physiol*, 262:R1137-R1144 (1992)), and this effect appears to be mediated, in part, by increased calcium sensitivity (Bark et al, *Hypertension*, 24:322-328 (1994)). Nitric oxide release from the cardiovascular regulatory center in the brain may also be involved in the central regulation of blood pressure, suggesting a role for neuronal NOS in the regulation of vascular tone (Cabrera et al, *Biochem Biophys Res Comm*, 206:77-81 (1995); Mattson et al, *Hypertension*, 28:297-303 (1996)).

Nitric oxide activates renin gene expression in the kidney, and is involved in the baroreceptor-mediated regulation of renin gene expression (Schricker et al, *Pflug Arch*, 428:261-268 (1994)). The dependence of blood pressure on salt intake appears to depend on NO, and NO deficiency states are associated with salt-sensitivity (Tolins et al, *Kidney Internat*, 46:230-236 (1994)). Selective inhibition of iNOS in Dahl R rats has been shown to lead to salt-sensitivity and to the development of salt-dependent hypertension similar to Dahl S rats (Rudd et al, *Am J Physiol*, 277: H732-H739 (1999)). In addition, mice deficient in iNOS (iNOS gene eliminated by targeted disruption) may develop hypertension in response to salt feeding (Rudd et al, *Circulation*, 98:1A (1998)).

Nitric oxide also affects myocardial contractility, and does so both by mediating muscarinic-cholinergic slowing of the heart rate and the contractile response to beta-adrenergic stimulation (Balligand et al, *Proc Nat'l Acad Sci USA*, 90:347-351 (1993)). This latter effect appears to be mediated *in vivo* through the vagus nerve (Hare et al, *J Clin Invest*, 95:360-366 (1995)).

In both vascular smooth muscle cells and cardiomyocytes, NO inhibits cellular proliferation and limits the proliferative response to growth-promoting substances (Garg et al, *J Clin Invest*, 83:1774-1777 (1986)). Left ventricular hypertrophy tends to occur in adult hearts with inadequate capillary proliferation, and this may account for the microvascular ischemia noted in patients with hypertrophy. Capillary proliferation is generally held to be a rare event in normal adult mammalian hearts. However, recent data from a hypertensive rat model, in which left ventricular hypertrophy commonly occurs, show that treatment with a low-dose of an angiotensin-converting enzyme inhibitor insufficient to prevent hypertension and left ventricular hypertrophy can, nonetheless, evoke capillary angiogenesis. Compared with untreated controls, treatment with the angiotensin converting enzyme inhibitor increased myocardial capillary proliferation (Unger et

al, *Hypertension*, 20:478-482 (1992)), and this effect was believed to be a consequence of inhibiting the degradation and potentiating the action of bradykinin. Bradykinin increases myocardial blood flow by inducing release of NO from microvascular endothelial cells, and increased blood flow is a powerful stimulus for capillary proliferation (Mall et al, *Bas Res Cardiol*, 85:531-540 (1990)).

5 Normal metabolic processes in vascular cells are associated with the generation of reactive oxygen intermediates that must be neutralized to limit oxidative damage and cellular dysfunction. In the setting of common cardiovascular disorders or in the presence of common risk factors for atherothrombotic disease, reactive oxygen species (ROS) are generated in abundance, and their rate of  
10 synthesis and flux typically exceeds the capacity of endogenous antioxidant mechanisms. Hypercholesterolemia, hyperglycemia (Keaney et al, *Circulation*, 99:189-191 (1999)), cigarette smoking, hyperhomocysteinemia, hypertension, and frank atherosclerosis are all accompanied by an increase in plasma and tissue ROS  
15 generation. Superoxide anion, hydrogen peroxide, hydroxyl radical, peroxynitrite, and lipid peroxides all increase in these settings. What remains unknown is whether or not the increase in ROS in these disorder is a primary event, a secondary consequence of the underlying process, or both.

Endogenous antioxidants important for the neutralization (i.e., reduction) of  
20 ROS can be categorized into two groups: small-molecule antioxidants and antioxidant enzymes. The former group comprises molecules such as GSH, NADPH,  $\alpha$ -tocopherol, vitamin C, and ubiquinol-10; while the latter group comprises the superoxide dismutases, catalase, and glutathione peroxidases. Deficiencies in several of these molecular species have been shown to lead to  
25 increased steady-state levels of ROS and vascular dysfunction, including increased platelet activation, arterial thrombosis (Freedman et al, *J Clin Invest*, 97:979-987 (1996); Freedman et al, *Circulation*, 98:1481-1486 (1998)), and reduced production of platelet-derived NO (Kenet et al, *Arterio Thromb Vasc Biol*, 19(8): 2017-2023 (1999)), which is important for limiting expansion of a platelet thrombus (Freedman et al,  
30 *Circ Res*, 84:1416-142 (1999)).

ROS generation accompanies the vascular dysfunction associated with several models of atherothrombotic and hypertensive vascular diseases. Hyperhomocysteinemic mice (i.e., cystathionine  $\beta$ -synthase knock-out mice) (Eberhardt et al,

5 *Circulation*, 98:144 (1998)), cellular glutathione peroxidase-deficient mice (i.e., cellular glutathione peroxidase knock-out mice), and salt-induced hypertensive rats (i.e., salt-fed Dahl S rats) (Trollet et al, *Circulation*, 98:1-725 (1998)) all manifest increased vascular ROS, and this increase in ROS is accompanied by reduced NO bioactivity through oxidative inactivation. Endothelial function and NO availability can be improved by improving antioxidant status with a cysteine precursor (Vita et al, *J Clin Invest*, 101:1408-1414 (1998)). In addition,  $\alpha$ -tocopherol leads to platelet inhibition (Freedman et al, *Circulation*, 94:2434-2440 (1996)) as one mechanism of its atherothrombotic benefit (Stephens et al, *Lancet*, 347:781-786 (1996)). The present inventors have also discovered that salt-loading salt-sensitive individuals (Dahl S rats) leads to an approximate 5-fold increase in plasma  $F_2$ -isoprostanes (8-epi-prostaglandin  $F_{2a}$ ), and this increase precedes the development of florid hypertension. These data all support the role of oxidant stress in the genesis or evolution of vascular dysfunction and disease, and the importance of antioxidant mechanisms in preventing this pathobiology, particularly with regard to African Americans.

In support of the mechanisms illustrated above, minimum forearm vascular resistance is significantly higher among normotensive blacks than whites (Bassett et al, *Am J Hypertension*, 5:781-786 (1992)), and forearm blood-flow responses to isoproterenol are markedly attenuated in normotensive blacks, suggesting a blunted  $\beta_2$ -vasodilator response in these individuals (Lang et al, *N Engl J Med*, 333:155-160 (1995)). Blacks tend to have greater left ventricular mass than whites for any given level of blood pressure (Koren et al, *Am J Hypertension*, 6:815-823 (1993); Chaturvedi et al, *J Am Coll Cardiol*, 24:1499-1505 (1994)). While not quantitated in any necropsy study, this response is likely to be accompanied by inadequate capillary angiogenesis which, in turn, may account for the diastolic dysfunction and the microvascular ischemia observed in blacks. Interestingly, blacks have been observed to have low levels of urinary kallikrein (Zinner et al, *Am J Epidemiol*, 104:124-132 (1976); Levy et al, *J Clin Invest*, 60:129-138 (1977)), the enzyme responsible for the generation of bradykinin from high-molecular-weight kininogen. Thus, were a similar abnormality in bradykinin and bradykinin-mediated NO production to exist in the coronary vasculature, attenuated blood flow responses may result that would limit capillary angiogenic responses and prevent the endothelial proliferative effects



of locally derived NO.

As discovered and described herein, African Americans have a unique vascular diathesis that may serve as the basis for clinically important cardiovascular syndromes. For example, differences in the outcome of left ventricular dysfunction may be a consequence of the enhanced (perhaps salt-dependent) increase in oxidant stress coupled with microvascular endothelial dysfunction and an inadequately vascularized, hypertrophied left ventricle. This constellation of pathophysiological abnormalities may provide the substrate for the important differences in outcome between blacks and whites with left ventricular dysfunction (Dreis et al, *N Engl J Med*, 340:609-616 (1999)). In addition, these observations and their clinical consequences suggest that blacks with abnormal endothelial function and nitric oxide insufficiency states would derive direct and, perhaps, disproportionate clinical benefit from enhancing nitric oxide in the vasculature, either by improving endothelial function, providing exogenous nitric oxide donors, or both.

In view of the above, the present invention provides methods of treating and/or preventing vascular diseases characterized by nitric oxide (NO) insufficiency by administering a therapeutically effective amount of at least one antioxidant or a pharmaceutically acceptable salt thereof, and at least one of isosorbide dinitrate and isosorbide mononitrate, and, optionally, at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, and/or at least one compound used to treat cardiovascular diseases. For example, the patient can be administered an antioxidant and isosorbide dinitrate, or the patient can be administered an antioxidant and isosorbide mononitrate, or the patient can be administered an antioxidant, isosorbide dinitrate and isosorbide mononitrate. The vascular diseases characterized by NO insufficiency include, for example, hypertension (e.g., low-renin hypertension; salt-sensitive hypertension; low-renin, salt-sensitive hypertension; primary pulmonary hypertension; thromboembolic pulmonary hypertension; pregnancy-induced hypertension; renovascular hypertension), heart failure (e.g., microvascular cardiac ischemia), and left ventricular hypertrophy with disproportionate microvascularization, (i.e., inadequate vascularity) or diastolic dysfunction.

Another aspect of the present invention provides methods for treating and/or preventing vascular diseases characterized by nitric oxide insufficiency by administering to a patient a therapeutically effective amount of at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist and/or nitrosated renin inhibitor, and, optionally, at least one antioxidant and/or at least one compound used to treat cardiovascular diseases.

The present invention also provides methods of preventing and treating Raynaud's syndrome by administering a therapeutically effective amount of at least one antioxidant or a pharmaceutically acceptable salt thereof, and at least one of isosorbide dinitrate and isosorbide mononitrate, and, optionally, at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist and/or nitrosated renin inhibitor. For example, the patient can be administered an antioxidant and isosorbide dinitrate, or the patient can be administered an antioxidant and isosorbide mononitrate, or the patient can be administered an antioxidant, isosorbide dinitrate and isosorbide mononitrate. The antioxidant and isosorbide dinitrate or isosorbide mononitrate can be administered separately or as components of the same composition. Raynaud's syndrome is a condition that causes a loss of blood flow to the fingers, toes, nose and/or ears. The affected area turns white from the lack of circulation, then blue and cold, and finally numb. The affected area may also turn red, and may throb, tingle or swell.

Another aspect of the present invention provides novel transdermal patches comprising a therapeutically effective amount of at least one antioxidant and at least one of isosorbide dinitrate and isosorbide mononitrate and/or at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, and mixtures thereof.

In the present invention, the antioxidants include small-molecule antioxidants and antioxidant enzymes. Suitable small-molecule antioxidants include, but are not limited to, hydralazine compounds, glutathione, vitamin C, vitamin E, cysteine, N-

acetyl-cysteine,  $\beta$ -carotene, ubiquinone, ubiquinol-10, tocopherols, coenzyme Q, and the like. Suitable antioxidant enzymes include, but are not limited to, superoxide dismutase, catalase, glutathione peroxidase, and the like. Suitable antioxidants are described more fully in the literature, such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Twelfth Edition, Version 12:1, 1996; and on STN Express, file phar and file reg.

The preferred antioxidant is a hydralazine compound that is preferably administered in the form of a pharmaceutically acceptable salt; more preferably in the form of hydralazine hydrochloride. Hydralazine hydrochloride is commercially available from, for example, Lederle Standard Products (Pearl River, NY), and Par Pharmaceuticals Inc. (Spring Valley, NY).

Isosorbide dinitrate is commercially available, for example, under the trade names DILATRATE®-SR (Schwarz Pharma, Milwaukee, WI); ISORDIL® and ISORDILR TITRADOSE® (Wyeth Laboratories Inc., Philadelphia, PA); and SORBITRATE® (Zeneca Pharmaceuticals, Wilmington, DE).

Isosorbide mononitrate is commercially available, for example, under the trade names IMDUR® (A. B. Astra, Sweden); MONOKET® (Schwarz Pharma, Milwaukee, WI); and ISMO® (Wyeth-Ayerst company, Philadelphia, PA).

The nitrosated angiotensin-converting enzyme inhibitors, nitrosated beta-adrenergic blockers, nitrosated calcium channel blockers, nitrosated endothelin antagonists, nitrosated angiotensin II receptor antagonists and nitrosated renin inhibitors of the present invention include any known angiotensin-converting enzyme inhibitors, beta-adrenergic blockers, calcium channel blockers, endothelin antagonists, angiotensin II receptor antagonists and renin inhibitors that have been nitrosated through one or more sites such as oxygen (hydroxyl condensation), sulfur (sulfhydryl condensation) and/or nitrogen. The nitrosated compounds of the present invention can be prepared using conventional methods known to one skilled in the art. For example, known methods for nitrosating compounds are described in U.S. Patent Nos. 5,380,758 and 5,703,073; WO 97/27749; WO 98/19672; and Oae et al, *Org. Prep. Proc. Int.*, 15(3):165-198 (1983), the disclosures of each of which are incorporated by reference herein in their entirety. WO 98/21193 discloses nitrosated ACE inhibitors and nitrosated beta-adrenergic blockers, the disclosure of

which is incorporated by reference herein in its entirety. WO 99/00361 discloses nitrate salts of ACE inhibitors, the disclosure of which is incorporated by reference herein in its entirety.

5        Suitable angiotensin-converting enzyme inhibitors, include, but are not limited to, alacepril, benazepril, captopril, ceranapril, cilazapril, delapril, duinapril, enalapril, enalaprilat, fosinopril, imidapril, lisinopril, moveltipril, pentopril, perindopril, quinapril, ramipril, rentipril, spirapril, temocapril, trandolapril, zofenopril, and the like. Suitable angiotensin-converting enzyme inhibitors are described more fully in the literature, such as in Goodman and Gilman, The  
10    Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Twelfth Edition, Version 12:1, 1996; and on STN Express, file phar and file registry.

      Suitable beta-adrenergic blockers, include, but are not limited to, amosulalol, atenolol, betaxolol, bethanidine, bevantolol, bisoprolol, bopindolol, bufuralol,  
15    bunitrolol, bupranolol, butafilolol, carazolol, carteolol, carvedilol, celiprolol, cetamolol, dilevalol, epanolol, indenolol, mepindolol, metipranolol, metoprolol, moprolol, nadolol, nebivolol, nipradilol, penbutolol, pindolol, pronethalol, propranolol, sotalol, sulfinalol, talinolol, tertatolol, tilisolol, timolol, toliprolol, and the like. Suitable beta-adrenergic blockers are described more fully in the literature,  
20    such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Twelfth Edition, Version 12:1, 1996; and on STN Express, file phar and file registry.

      Suitable calcium channel blockers, include, but are not limited to, amlodipine, aranidipine, barnidipine, benidipine, cilnidipine, clentiazem, diltiazem, efonidipine,  
25    fantofarone, felodipine, isradipine, lacidipine, lercanidipine, manidipine, mibefradil, nicardipine, nifedipine, nilvadipine, nisoldipine, nitrendipine, semotiadil, veraparmil, and the like. Suitable calcium channel blockers are described more fully in the literature, such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM,  
30    Twelfth Edition, Version 12:1, 1996; and on STN Express, file phar and file registry.

      Suitable endothelin antagonists, include, but are not limited to, bosentan, sulfonamide endothelin antagonists, BQ-123, SQ 28608, and the like. Suitable endothelin antagonists are described more fully in the literature, such as in

Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Twelfth Edition, Version 12:1, 1996; and on STN Express, file phar and file registry.

5        Suitable angiotensin II receptor antagonists, include, but are not limited to, ciclosidomine, eprosartan, furosemide, irbesartan, losartan, saralasin, valsartan, and the like. Suitable angiotensin II receptor antagonists are described more fully in the literature, such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Twelfth Edition, Version 12:1, 1996; and on STN Express, file phar and file registry.

10        Suitable renin inhibitors, include, but are not limited to, enalkrein, RO 42-5892, A 65317, CP 80794, ES 1005, ES 8891, SQ 34017, and the like). Suitable renin inhibitors are described more fully in the literature, such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Twelfth Edition, Version 12:1, 1996; and on  
15        STN Express, file phar and file registry.

      In the methods of the invention, the antioxidant and at least one of isosorbide dinitrate and isosorbide mononitrate, and, optionally, at least one nitrosated compound, and/or compound used to treat cardiovascular diseases can be administered as separate components or as components of the same composition.  
20        When the antioxidant and at least one of isosorbide dinitrate and isosorbide mononitrate are administered as separate components for the treatment of vascular diseases characterized by NO insufficiency or Raynaud's syndrome, they are preferably administered to the patient at about the same time. "About the same time" means that within about thirty minutes of administering one compound (e.g.,  
25        antioxidant or isosorbide dinitrate or isosorbide mononitrate) to the patient, the other compound (e.g., isosorbide dinitrate or isosorbide mononitrate or antioxidant) is administered to the patient. "About the same time" also includes simultaneous administration of the compounds.

      In addition to the administration of the combination of the antioxidant and isosorbide dinitrate or isosorbide mononitrate for the treatment of vascular diseases  
30        characterized by NO insufficiency, the patients can receive digitalis such as digoxin and/or diuretics and/or at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker,

nitrosated endothelin antagonist, angiotensin II receptor antagonist, nitrosated renin inhibitor, and/or at least one compound used to treat cardiovascular diseases.

5 The digoxin is preferably administered orally to achieve a steady state blood serum concentration of at least about 0.7 nanograms per ml to about 2.0 nanograms per ml. The diuretic is administered, preferably orally, to manage edema. Suitable  
10 diuretics include, but are not limited to, thiazides (such as, for example, chlorothiazide, hydrochlorothiazide); ethacrynic acid, furosemide, spironalactone, triamterene or mixtures thereof. Depending on the diuretic used, potassium may also be administered to the patient in order to optimize the fluid balance while  
15 avoiding hypokalemic alkalosis. The administration of potassium can be in the form of potassium chloride or by the daily ingestion of foods with high potassium content such as, for example, bananas, orange juice, and the like. The method of administration of these compounds is described in further detail in U.S. Patent No. 4,868,179, the disclosure of which is incorporated by reference herein in its entirety.

20 The compounds and compositions of the invention can be administered by any available and effective delivery system including, but not limited to, orally, buccally, parenterally, by inhalation spray, topically (including transdermally), or rectally in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. The preferred methods of administration are by oral administration or topical application (transdermal application).

Topical administration can also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. Dosage forms for topical administration of the compounds and compositions can include creams, sprays,  
25 lotions, gels, ointments, and the like. In such dosage forms, the compositions of the invention can be mixed to form white, smooth, homogeneous, opaque cream or lotion with, for example, benzyl alcohol 1% or 2% (wt/wt) as a preservative, emulsifying wax, glycerin, isopropyl palmitate, lactic acid, purified water and sorbitol solution. In addition, the compositions can contain polyethylene glycol 400.  
30 They can be mixed to form ointments with, for example, benzyl alcohol 2% (wt/wt) as preservative, white petrolatum, emulsifying wax, and tenox II (butylated hydroxyanisole, propyl gallate, citric acid, propylene glycol). Woven pads or rolls of bandaging material, e.g., gauze, can be impregnated with the compositions in

solution, lotion, cream, ointment or other such form can also be used for topical application.

The compositions can also be applied topically using a transdermal system, such as one of an acrylic-based polymer adhesive with a resinous crosslinking agent impregnated with the composition and laminated to an impermeable backing. In a preferred embodiment, the compositions of the present invention are administered in the form of a transdermal patch, more preferably in the form of a sustained-release transdermal patch. The transdermal patches of the present invention can include any conventional form such as, for example, adhesive matrix, polymeric matrix, reservoir patch, matrix or monolithic-type laminated structure, and are generally comprised of one or more backing layers, adhesives, penetration enhancers, an optional rate controlling membrane and a release liner which is removed to expose the adhesives prior to application. Polymeric matrix patches also comprise a polymeric-matrix forming material. Suitable transdermal patches are described in more detail in, for example, U. S. Patent Nos. 5,262,165, 5,948,433, 6,010,715 and 6,071,531, the disclosure of each of which are incorporated herein in their entirety.

Solid dosage forms for oral administration can include capsules, sustained-release capsules, tablets, chewable tablets, sublingual tablets, effervescent tablets, pills, powders, granules and gels. In such solid dosage forms, the active compounds can be admixed with at least one inert diluent, such as sucrose, lactose or starch. Such dosage forms can also comprise, as is normal practice, additional substances other than inert diluents, e.g., lubricating agents, such as magnesium stearate. In the case of capsules, tablets, effervescent tablets, and pills, the dosage forms can also comprise buffering agents. Soft gelatin capsules can be prepared to contain a mixture of the active compound or composition and vegetable oil. Hard gelatin capsules can contain granules of the active compound in combination with a solid, pulverulent carrier, such as lactose, saccharose, sorbitol, mannitol, potato starch, corn starch, amylopectin, or cellulose derivatives of gelatin. Tablets and pills can be prepared with enteric coatings.

Liquid dosage forms for oral administration can include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions can also

comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

Suppositories for rectal administration of the compounds or compositions can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols which are solid at room temperature but liquid at  
5      rectal temperature, such that they will melt in the rectum and release the drug.

The term parenteral includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques. Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions can be formulated  
10      according to the known art using suitable dispersing agents, wetting agents and/or suspending agents. The sterile injectable preparation can also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be used are water, Ringer's solution, and isotonic sodium chloride  
15      solution. Sterile fixed oils are also conventionally used as a solvent or suspending medium.

The compounds and compositions of the invention will typically be administered in a pharmaceutical composition comprising one or more carriers or excipients. Examples of suitable carriers include, for example, water, silicone,  
20      waxes, petroleum jelly, polyethylene glycols, propylene glycols, liposomes, sugars, salt solutions, alcohol, vegetable oils, gelatins, lactose, amylose, magnesium stearate, talc, surfactants, silicic acids, viscous paraffins, perfume oils, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethylcelluloses, polyvinyl-pyrrolidones, and the like. The pharmaceutical  
25      preparations can be sterilized and if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously react with the active compounds. For topical application, the compositions can also include one or more permeation enhancers  
30      including, for example, dimethylsulfoxide (DMSO), dimethyl formamide (DMF), N,N-dimethylacetamide (DMA), decylmethylsulfoxide (C10MSO), polyethylene glycol monolaurate (PEGML), glycerol monolaurate, lecithin, 1-substituted azacycloheptan-2-ones, particularly 1-N-dodecylcyclazacycoheptan-2-ones (available



under the trademark AZONE from Nelson Research & Development Co., Irvine CA), alcohols and the like. For parenteral application, particularly suitable vehicles consist of solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants. Aqueous suspensions can contain substances which  
5 increase the viscosity of the suspension and include, for example, sodium carboxymethyl cellulose, sorbitol and/or dextran. Optionally, the suspension can contain stabilizers. The compositions, if desired, can also contain minor amounts of wetting agents, emulsifying agents and/or pH buffering agents.

Various delivery systems are known and can be used to administer the  
10 compounds or compositions of the present invention, including, for example, encapsulation in liposomes, microbubbles, emulsions, microparticles, microcapsules and the like. The required dosage can be administered as a single unit or in a sustained release form.

While individual needs may vary, determination of optimal ranges for  
15 effective amounts of the compounds and/or compositions is within the skill of the art and can be determined by standard clinical techniques, including reference to Goodman and Gilman, supra; The Physician's Desk Reference, Medical Economics Company, Inc., Oradell, N.J., 1995; and Drug Facts and Comparisons, Inc., St. Louis, MO, 1993. Generally, the dosage required to provide an effective amount of the  
20 compounds and compositions, which can be adjusted by one of ordinary skill in the art, will vary depending on the age, health, physical condition, sex, diet, weight, extent of the dysfunction of the recipient, frequency of treatment and the nature and scope of the dysfunction or disease, medical condition of the patient, the route of administration, pharmacological considerations such as the activity, efficacy,  
25 pharmacokinetic and toxicology profiles of the particular compound used, whether a drug delivery system is used, and whether the compound is administered as part of a drug combination.

The compounds and compositions of the present invention can be formulated  
as pharmaceutically acceptable salts. Pharmaceutically acceptable salts include, for  
30 example, alkali metal salts and addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids include, but are not

limited to, hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid and the like. Appropriate organic acids include, but are not limited to, aliphatic, cycloaliphatic, aromatic, heterocyclic, carboxylic and sulfonic classes of organic acids, such as, for example, formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicylic, p-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, stearic, algenic,  $\beta$ -hydroxybutyric, cyclohexylaminosulfonic, galactaric and galacturonic acid and the like. Suitable pharmaceutically-acceptable base addition salts include, but are not limited to, metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from primary, secondary and tertiary amines, cyclic amines, N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine and the like. All of these salts may be prepared by conventional means from the corresponding compound by reacting, for example, the appropriate acid or base with the compound.

In preferred embodiments, the hydralazine is administered in an amount of about 30 milligrams per day to about 300 milligrams per day; the isosorbide dinitrate is administered in an amount of about 20 milligrams per day to about 200 milligrams per day; and the isosorbide mononitrate is administered in an amount of about 10 milligrams per day to about 120 milligrams per day. The preferred amounts of hydralazine and/or isosorbide dinitrate or isosorbide mononitrate can be administered as a single dose once a day; in multiple doses several times throughout the day; or in a sustained-release formulation.

The present invention also provides pharmaceutical kits comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compounds and/or compositions. Such kits can also include, for example, other compounds and/or compositions (e.g., diuretics, digoxin, nitrosated compounds, compounds used to treat cardiovascular diseases and the like), a device(s) for administering the compounds and/or compositions, and written instructions in a

form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which instructions can also reflect approval by the agency of manufacture, use or sale for human administration.

### EXAMPLES

5       The following examples are for purposes of illustration only, and are not intended to limit the scope of the specification or claims.

#### Example 1

10       As described herein, NO deficiency is a central pathophysiologic mechanism for the black vascular diathesis. To examine this issue in forearm microvessels, the vasodilator responses to intra-arterial infusions of methacholine, sodium nitroprusside, and verapamil were examined using venous occlusion plethysmography in 36 white and black hypertensive patients. These patients had no other coronary factors, such as smoking, diabetes mellitus, or hypercholesterolemia, and the two groups were matched in terms of age, gender, 15 lipid levels, blood pressure, and anti-hypertensive treatment. Similar to previous reports (Lang et al, *N Engl J Med*, 333:155-160 (1995); Panza et al, *N Engl J Med*, 323:22-27 (1990)), the dilator response to methacholine, but not nitroprusside, was significantly reduced in these hypertensive patients compared to age-matched normotensive controls (Sherman et al, *Circulation*, 98:1-376 (1998)). Regarding racial 20 differences, as shown in Fig. 2, vasodilation in response to methacholine was markedly worse in the black hypertensive patients compared to white hypertensive patients. There were no racial differences in the responses to sodium nitroprusside or verapamil (data not shown), suggesting that this impairment of NO action in black hypertensives is at the endothelial level and that dysfunction of vascular 25 smooth muscle does not account for the impaired response. These findings have important implications for the pathogenesis of hypertension and myocardial ischemia.

#### Example 2

30       To investigate the issue of whether NO deficiency is a primary or secondary phenomenon in hypertensive black patients, 46 normotensive white and black patients who were matched for age and gender were compared. As shown in Fig. 3B, the vasodilator responses to methacholine were not significantly different in black and white normotensive patients. However, the response to sodium

nitroprusside (Fig. 3A) was significantly lower in the black normotensive patients. This finding shows that there is an impairment in the vasodilator response exogenous source of NO in black patients even prior to the development of hypertension, an observation that would be consistent with a primary rather than secondary role in the black vascular diathesis. These findings are consistent with recently published results by other investigators (Lang et al, *N Engl J Med*, 333:155-160 (1995); Cardillo et al, *Hypertension* 31:1235-1239 (1998)).

### Example 3

Black patients have a preponderance of salt-sensitive hypertension, and the data described herein show that NO deficiency is a pathogenic mechanism of salt-sensitive hypertension in experimental models (Rudd et al, *Am J Physiol*, 277: H732-H739 (1999)). To explore the relation between salt-sensitivity and endothelium-derived NO action in black patients prior to the development of hypertension, the blood pressure response to salt-loading and salt-deprivation using an established inpatient protocol (Weinberger et al, *Hypertension*, 8:II-127-II-134 (1986)) in a group of normotensive black patients was assessed. Briefly, blood pressure was continuously monitored non-invasively during a salt load (sodium 458 mEq over 24 hours) and during a period of salt depletion (furosemide treatment and sodium intake 10 mEq over 24 hours). Patients were considered to be salt-sensitive if mean blood pressure was at least 10 mm Hg higher during the salt loading period (Weinberger et al, *Hypertension*, 8:II-127-II-134 (1986)). As shown in Figs. 4A-B, there were trends for impaired vasodilator responses to both methacholine (Fig. 4A) and sodium nitroprusside (Fig. 4B). The vasodilator responses to verapamil were equivalent in salt-sensitive and salt-resistant patients (data not shown). These data show impaired NO action in the microvasculature of salt sensitive individuals prior to the development of hypertension.

### Example 4

The effects of hypertension and race on conduit vessel function were examined using a well-established brachial ultrasound technique (Vita et al, Lanzer & Lipton, Eds., Diagnostics of Vascular Diseases: Principles and Technology. Berlin:Springer-Verlag, pp.249-259 (1996)) in 370 patients (178 black, 192 white). As shown in Table 1, there were no significant differences in flow-mediated dilation or nitroglycerin-mediated dilation according to race. However, hypertension was

associated with a highly significant reduction in both flow-mediated dilation and nitroglycerin-mediated dilation. Further, systolic blood pressure was inversely correlated with flow-mediated dilation ( $r = -0.30$ ,  $P < 0.001$ ) and nitroglycerin-mediated dilation ( $r = -0.33$ ,  $P < 0.001$ ). Diastolic pressure correlated to a similar extent with vascular function. By multiple linear regression analysis, vessel size and systolic blood pressure were the only independent predictors of flow-mediated dilation in this sizable group of patients. Thus, both black and white patients with hypertension demonstrate a significant impairment of NO action in conduit arteries (Gokce et al, *Circulation*, 99(25) 3234-3240 (1999)).

Table 1: Conduit Vasomotor Function By Race

	White Patients	Black Patients	P
FMD (%)			
Normotensive	11.5 ± 5.9	13.3 ± 7.2	NS
Hypertensive	9.5 ± 5.3*	8.8 ± 6.2*	NS
NTG-response (%)			
Normotensive	18.1 ± 8.1	20.6 ± 8.4	NS
Hypertensive	14.6 ± 6.2*	16.4 ± 7.1*	NS

Data are mean ± SD. \*p < 0.001 compared to race matched normotensives.

Each of the above examples demonstrate that NO action is impaired in the microvasculature of black hypertensive patients to a greater extent than in white hypertensive patients. There is a suggestion that this abnormality may precede the development of hypertension, particularly in salt-sensitive individuals, consistent with a pathogenic role. In conduit vessels, a marked impairment of NO action that may contribute to ischemic heart disease and stroke has been demonstrated. This abnormality appears to be independent of race, a finding that is consistent with possibility that the impairment is a consequence of blood pressure elevation of any cause. Since African Americans have a greater incidence of hypertension and a preponderance of salt-sensitive hypertension, these findings lend further support to the hypothesis that NO insufficiency contributes to the pathogenesis of ischemic heart disease in this population.

The disclosure of each patent, patent application and publication cited herein

is hereby incorporated by reference herein in its entirety.

Although the invention has been set forth in detail, one skilled in the art will appreciate that numerous changes and modifications can be made to the invention without departing from the spirit and scope thereof.

## CLAIMS

What is claimed is:

1. A method of treating and/or preventing a vascular disease characterized by nitric oxide insufficiency in a patient comprising administering to the patient a therapeutically effective amount of at least one antioxidant, or a pharmaceutically acceptable salt thereof, and at least one of isosorbide dinitrate and isosorbide mononitrate.
2. The method of claim 1, wherein the vascular disease characterized by nitric oxide insufficiency is low-renin hypertension; salt-sensitive hypertension; low-renin, salt-sensitive hypertension; primary pulmonary hypertension; thromboembolic pulmonary hypertension; pregnancy-induced hypertension; renovascular hypertension; heart failure; microvascular cardiac ischemia; left ventricular hypertrophy with disproportionate microvascularization or diastolic dysfunction.
3. The method of claim 1, further comprising administering a pharmaceutically acceptable carrier.
4. The method of claim 1, wherein the antioxidant is a small-molecule antioxidant, or a pharmaceutically acceptable salt thereof, or an antioxidant enzyme.
5. The method of claim 4, wherein the small-molecule antioxidant is a hydralazine compound, a glutathione, a vitamin C, a vitamin E, a cysteine, a N-acetyl-cysteine, a  $\beta$ -carotene, an ubiquinone, an ubiquinol-10, a tocopherol, a coenzyme Q, or a mixture thereof.
6. The method of claim 4, wherein the antioxidant enzyme is a superoxide dismutase, a catalase, a glutathione peroxidase, or a mixture thereof.
7. The method of claim 5, wherein the hydralazine compound is budralazine, cadralazine, dihydralazine, endralazine, hydralazine, pildralazine or todralazine or a pharmaceutically acceptable salt thereof.
8. The method of claim 7, wherein the hydralazine compound is hydralazine hydrochloride.
9. The method of claim 8, wherein the hydralazine hydrochloride is administered in an amount of about 30 milligrams per day to about 300 milligrams per day.
10. The method of claim 1, wherein the isosorbide dinitrate is

administered in an amount of about 20 milligrams per day to about 200 milligrams per day.

11. The method of claim 1, wherein the isosorbide mononitrate is administered in an amount of about 10 milligrams per day to about 120 milligrams per day.

12. The method of claim 1, wherein the at least one antioxidant and the at least one of isosorbide dinitrate and isosorbide mononitrate are administered orally.

13. The method of claim 12, wherein the at least one antioxidant and the at least one of isosorbide dinitrate and isosorbide mononitrate are orally administered in the form of a solid dose.

14. The method of claim 13, wherein the solid dose is in the form of a tablet or capsule.

15. The method of claim 14, wherein the capsule is in the form of a sustained release capsule

16. The method of claim 14, wherein the tablet is in the form of a sublingual tablet.

17. The method of claim 14, wherein the tablet is in the form of a chewable tablet.

18. The method of claim 1, wherein the at least one antioxidant and the at least one of isosorbide dinitrate and isosorbide mononitrate are administered as components of the same composition.

19. The method of claim 1, wherein the at least one antioxidant and the at least one of isosorbide dinitrate and isosorbide mononitrate are administered to the patient as separate components.

20. The method of claim 19, wherein the at least one antioxidant and the at least one isosorbide dinitrate and isosorbide mononitrate are administered to the patient as separate components at about the same time.

21. The method of claim 1, wherein the at least one antioxidant and the at least one of isosorbide dinitrate and isosorbide mononitrate are administered transdermally.

22. The method of claim 21, wherein the transdermal application is in the form of a sustained-release patch.

23. The method of claim 1, further comprising administering a digitalis.



24. The method of claim 23, wherein the digitalis is digoxin
25. The method of claim 24, wherein the digoxin is administered in an amount to achieve a blood serum concentration of at least about 0.7 nanograms per milliliter to about 2.0 nanograms per milliliter.
- 5 26. The method of claim 1 further comprising administering a therapeutically effective edema managing amount of a diuretic compound.
27. The method of claim 26, wherein the diuretic compound is a thiazide, ethacrynic acid, a furosemide, a spiranolactone, a triamterene, or a mixture thereof.
- 10 28. The method of claim 26, further comprising administering a therapeutically effective amount of potassium.
29. The method of claim 28, wherein the potassium is administered in the form of potassium chloride or by the daily ingestion of foods with high potassium content.
- 15 30. The method of claim 1, further comprising administering at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, or a mixture thereof.
- 20 31. The method of claim 1, further comprising administering at least one compound used to treat cardiovascular diseases, or a pharmaceutically acceptable salt thereof.
- 25 32. The method of claim 31, wherein the at least one compound used to treat cardiovascular diseases is an angiotensin-converting enzyme inhibitor, a beta-adrenergic blocker, a cholesterol reducer, a calcium channel blocker, an angiotensin II receptor antagonist, an endothelin antagonist, a renin inhibitor, or a mixture thereof.
- 30 33. The method of claim 1, comprising administering to the patient a therapeutically effective amount of hydralazine hydrochloride and isosorbide dinitrate.
34. The method of claim 1, comprising administering to the patient a therapeutically effective amount of hydralazine hydrochloride and isosorbide mononitrate.
35. A method of treating Raynaud's syndrome in a patient comprising

administering to the patient a therapeutically effective amount of at least one antioxidant, or a pharmaceutically acceptable salt thereof, and at least one of isosorbide dinitrate and isosorbide mononitrate.

5 36. The method of claim 35, further comprising administering a pharmaceutically acceptable carrier.

37. The method of claim 35, wherein the antioxidant is a small-molecule antioxidant, or a pharmaceutically acceptable salt thereof, or an antioxidant enzyme.

10 38. The method of claim 37, wherein the small-molecule antioxidant is a hydralazine compound, a glutathione, a vitamin C, a vitamin E, a cysteine, a N-acetyl-cysteine, a  $\beta$ -carotene, an ubiquinone, an ubiquinol-10, a tocopherol, a coenzyme Q, or a mixture thereof.

39. The method of claim 37, wherein the antioxidant enzyme is a superoxide dismutase, a catalase, a glutathione peroxidase, or a mixture thereof.

15 40. The method of claim 38, wherein the hydralazine compound is budralazine, cadralazine, dihydralazine, endralazine, hydralazine, pildralazine or todralazine or a pharmaceutically acceptable salt thereof.

41. The method of claim 40, wherein the hydralazine compound is hydralazine hydrochloride.

20 42. The method of claim 35, comprising administering to the patient a therapeutically effective amount of hydralazine hydrochloride and isosorbide dinitrate.

43. The method of claim 35, comprising administering to the patient a therapeutically effective amount of hydralazine hydrochloride and isosorbide mononitrate.

25 44. The method of claim 41, wherein the hydralazine hydrochloride is administered in an amount of about 20 milligrams per day to about 200 milligrams per day.

45. The method of claim 35, wherein the isosorbide dinitrate is administered in an amount of about 20 milligrams per day to about 200 milligrams per day.

30 46. The method of claim 35, wherein the isosorbide mononitrate is administered in an amount of about 10 milligrams per day to about 120 milligrams per day.

47. The method of claim 35, wherein the at least one antioxidant and the at least one of isosorbide dinitrate and isosorbide mononitrate are administered transdermally.

5 48. The method of claim 47, wherein the transdermal application is in the form of a sustained-release patch.

49. The method of claim 35, wherein the at least one antioxidant and the at least one of isosorbide dinitrate and isosorbide mononitrate are administered orally.

50. The method of claim 49, wherein the oral administration is in the form of a solid dose.

10 51. The method of claim 50, wherein the solid dose is a tablet or a capsule.

52. The method of claim 50, wherein the capsule is a sustained-release capsule.

53. The method of claim 51, wherein the tablet is a sublingual tablet or a chewable tablet.

15 54. The method of claim 35, wherein the at least one antioxidant and the at least one of isosorbide dinitrate and isosorbide mononitrate are administered as components of the same composition.

55. The method of claim 35, wherein the at least one antioxidant and the at least one of isosorbide dinitrate and isosorbide mononitrate are administered as separate components.

20 56. The method of claim 35, further comprising administering at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, or a mixture thereof.

25 57. A method of treating and/or preventing a vascular disease characterized by nitric oxide insufficiency in a patient comprising administering to the patient a therapeutically effective amount of at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, or a mixture thereof.

30 58. The method of claim 57, further comprising administering a pharmaceutically acceptable carrier.

59. The method of claim 57, wherein the at least one nitrosated

angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, or mixture thereof, are in the form of a pharmaceutically acceptable salt.

5           60. The method of claim 57, wherein the at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, or mixture thereof, are orally administered in the form of a solid dose.

10           61. The method of claim 60, wherein the solid dose is in the form of a tablet or capsule.

          62. The method of claim 61, wherein the capsule is in the form of a sustained release capsule

15           63. The method of claim 61, wherein the tablet is in the form of a sublingual tablet.

          64. The method of claim 61, wherein the tablet is in the form of a chewable tablet.

20           65. The method of claim 57, wherein the at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist, nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, or a mixture thereof, are administered transdermally.

          66. The method of claim 65, wherein the transdermal application is in the form of a sustained-release patch.

25           67. The method of claim 57, further comprising administering at least one antioxidant, or a pharmaceutically acceptable salt thereof.

          68. The method of claim 67, wherein the antioxidant is a small-molecule antioxidant, or a pharmaceutically acceptable salt thereof, or an antioxidant enzyme.

30           69. The method of claim 68, wherein the small-molecule antioxidant is a hydralazine compound, a glutathione, a vitamin C, a vitamin E, a cysteine, a N-acetyl-cysteine, a  $\beta$ -carotene, an ubiquinone, an ubiquinol-10, a tocopherol, a coenzyme Q, or a mixture thereof.

          70. The method of claim 68, wherein the antioxidant enzyme is a

superoxide dismutase, a catalase, a glutathione peroxidase, or a mixture thereof.

71. The method of claim 69, wherein the hydralazine compound is budralazine, cadralazine, dihydralazine, endralazine, hydralazine, pildralazine or todralazine, or a pharmaceutically acceptable salt thereof.

5 72. The method of claim 71, wherein the hydralazine compound is hydralazine hydrochloride.

73. The method of claim 57, further comprising administering at least one compound used to treat cardiovascular diseases, or a pharmaceutically acceptable salt thereof.

10 74. The method of claim 73, wherein the at least one compound used to treat cardiovascular diseases is an angiotensin-converting enzyme inhibitor, a beta-adrenergic blocker, a cholesterol reducer, a calcium channel blocker, an angiotensin II receptor antagonist, an endothelin antagonist, a renin inhibitor, or a mixture thereof.

15 75. A transdermal patch comprising a therapeutically effective amount of at least one antioxidant and at least one of isosorbide dinitrate and isosorbide mononitrate.

76. The transdermal patch of claim 75, further comprising a pharmaceutically acceptable carrier.

20 77. The transdermal patch of claim 75, wherein the antioxidant is a small-molecule antioxidant, or a pharmaceutically acceptable salt thereof, or an antioxidant enzyme.

25 78. The transdermal patch of claim 77, wherein the small-molecule antioxidant is a hydralazine compound, a glutathione, a vitamin C, a vitamin E, a cysteine, a N-acetyl-cysteine, a  $\beta$ -carotene, an ubiquinone, an ubiquinol-10, a tocopherol, a coenzyme Q, or a mixture thereof.

79. The transdermal patch of claim 77, wherein the antioxidant enzyme is superoxide dismutase, catalase, glutathione peroxidase, or a mixture thereof.

30 80. The transdermal patch of claim 78, wherein the at least one antioxidant is a hydralazine compound or a pharmaceutically acceptable salt thereof.

81. The transdermal patch of claim 80, wherein the at least one hydralazine compound is hydralazine hydrochloride.

82. The transdermal patch of claim 81, wherein the hydralazine

hydrochloride is present in an amount of about 30 milligrams to about 300 milligrams.

83. The transdermal patch of claim 75, wherein the isosorbide dinitrate is present in an amount of about 20 milligrams per day to about 200 milligrams per day.

84. The transdermal patch of claim 75, wherein the isosorbide mononitrate is administered in an amount of about 10 milligrams per day to about 120 milligrams per day.

85. The transdermal patch of claim 75, wherein the transdermal patch is a sustained-release transdermal patch.

86. The transdermal patch of claim 75, comprising a therapeutically effective amount of hydralazine hydrochloride and isosorbide dinitrate.

87. The transdermal patch of claim 75, comprising a therapeutically effective amount of hydralazine hydrochloride and isosorbide mononitrate.

88. The transdermal patch of claim 75, further comprising administering at least one nitrosated angiotensin-converting enzyme inhibitor, nitrosated beta-adrenergic blocker, nitrosated calcium channel blocker, nitrosated endothelin antagonist and nitrosated angiotensin II receptor antagonist, nitrosated renin inhibitor, or a mixture thereof.

89. A method of treating a vascular disease characterized by a nitric oxide insufficiency in a patient comprising administering to the patient the transdermal patch of claim 75.

90. The method of claim 89, wherein the vascular disease characterized by nitric oxide insufficiency is low-renin hypertension; salt-sensitive hypertension; low-renin, salt-sensitive hypertension; primary pulmonary hypertension; thromboembolic pulmonary hypertension; pregnancy-induced hypertension; renovascular hypertension; heart failure; microvascular cardiac ischemia; left ventricular hypertrophy with disproportionate microvascularization or diastolic dysfunction.

91. A method of treating Raynaud's syndrome in a patient comprising administering to the patient the transdermal patch of claim 75.

FIG. 1





FIG. 2



2/3

FIG. 3A




FIG. 3B






FIG. 4A





FIG. 4B



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/29528

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/535, 31/50, 31/495, 31/415, 31/355, 31/34, 31/19, 31/135; A01N 45/00  
US CL : 514/26, 233.8, 248, 398, 400, 401, 458, 474, 571, 647, 654, 953

According to International Patent Classification (IPC) or to both national classification and IPC.

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
U.S. : 514/26, 233.8, 248, 398, 400, 401, 458, 474, 571, 647, 654, 953

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Please See Continuation Sheet

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,627,191 A (BIRCH et al.) 06 May 1997(06.05.1997) see abstract, column 33, line 56 to column 34, line 36.	1-74 and 89-91



Further documents are listed in the continuation of Box C.



See patent family annex.

\* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent published on or after the international filing date
- "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

16 APR 2001

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer  
Marianne Cintas

Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/29528

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claim Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:  
Please See Continuation Sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☒

No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/29528

**BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING** This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claim(s) 1-74 and 89-91 drawn to a method of treating and/or preventing a vascular disease.

Group II, claim(s) 75-88, drawn to a transdermal patch.

The inventions listed as Groups I-II do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: A patch is not required to practice the invention of group I.

**Continuation of B. FIELDS SEARCHED Item3:** WEST 2.0, STN file MEDICINE, search terms: vascular, digoxin, digitalis, furosemide, triamterene, isosorbide dinitrate, or mononitrate, hydralazine, dihydralazine, bupralazine, cadralazine, endralazine, pildralazine, todralazine, cysteine, superoxide, dismutase, glutathione, beta carotene, Vitamin C, Vitamine E, tocopherol, ascorbic acid, angiotensin II, Reynaud, thiazide, ethacrynic acid